



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

tion of brain, meningitis, and membranous croup were less prevalent than in the preceding week.

Meningitis was reported present at 4, whooping cough at 14, diphtheria at 30, enteric fever at 45, pneumonia at 57, scarlet fever at 68, measles at 77, smallpox at 87, and phthisis pulmonalis at 227 places.

The Monthly Bulletin of Vital Statistics says :

The total number of deaths returned to the Department of State for the month of March was 3,406, which was only 19 more than the number returned for the preceding month. The death rate for March was 16.0 per 1,000 population, as compared with 17.1 for the preceding month. Important causes of death were as follows: Tuberculosis of lungs, 227; other forms of tuberculosis, 43; enteric fever, 57; diphtheria and croup, 38; scarlet fever, 31; measles, 27; whooping cough, 21; pneumonia, 384; influenza, 156; cancer, 149; accidents and violence, 199.

Deaths from pulmonary tuberculosis and enteric fever showed some increase, while diphtheria and croup, pneumonia and influenza caused considerably fewer deaths in March than in February.

By ages, there were 527 deaths of infants under 1 year of age; 185 deaths of children aged 1 to 4 years, inclusive; and 1,159 deaths of elderly persons 65 years and over.

There were 3 deaths from smallpox, 1 in Casnovia village, Muskegon County; 1 in Leonidas Township, St. Joseph County; and 1 in Evart Township, Osceola County.

TENNESSEE—Nashville.—Month of March, 1904. Estimated population, 82,947; white, 52,061; colored, 30,886. Total number of deaths, 184; white, 86; colored, 98, including enteric fever 1, measles 11, scarlet fever 2, and 34 from tuberculosis.

UTAH—Salt Lake City.—Month of March, 1904. Census population, 53,531. Total number of deaths, 117, including diphtheria 2, enteric fever 1, whooping cough 7, and 14 from tuberculosis.

Report of immigration at Baltimore for the week ended April 16, 1904.

OFFICE OF THE COMMISSIONER OF IMMIGRATION,
Baltimore, Md., April 16, 1904.

Number of alien immigrants who arrived at this port during the week ended April 16, 1904; also names of vessels and ports from which they came.

Date of arrival.	Vessel.	Where from.	Number of immigrants.
Apr. 13 16	Brandenburg.....	Bremen	1,039
	Frankfurt	do	941
	Total	1,980

LOUIS T. WEIS, *Commissioner.*